arXiv:1511.06932v3 [math.PR] 14 Sep 2016

First passage percolation on the exponential of two-dimensional branching random walk

Jian Ding* University of Chicago

Subhajit Goswami* University of Chicago

September 16, 2016

Abstract

We consider the branching random walk $\{\mathcal{R}_z^N : z \in V_N\}$ with Gaussian increments indexed over a two-dimensional box V_N of side length N, and we study the first passage percolation where each vertex is assigned weight $e^{\gamma \mathcal{R}_z^N}$ for $\gamma > 0$. We show that for $\gamma > 0$ sufficiently small but fixed, the expected FPP distance between the left and right boundaries is at most $O(N^{1-\gamma^2/10})$.

Key words and phrases. First passage percolation (FPP), Branching Random Walk (BRW), Gaussian free field (GFF), Liouville quantum gravity (LQG).

1 Introduction

For $n \in \mathbb{N}$ and $N = 2^n$, let $V_N \subseteq \mathbb{Z}^2$ be a box of side length N whose lower left corner is located at the origin. Let $\mathbb{BD}_j$ denote the collection of boxes of the form $([0, 2^j - 1] \cap \mathbb{Z})^2 + (i_1 2^j, i_2 2^j)$ where i_1, i_2 and j are nonnegative integers. For $v \in \mathbb{Z}^2$, let $\mathbb{BD}_j(v)$ be the (unique) box in $\mathbb{BD}_j$ that contains v. Let $\{a_B\}_{k \geq 0, B \in \mathbb{BD}_k}$ be a collection of i.i.d. standard Gaussian variables. We define a branching random walk (indexed over V_N) with Gaussian increments by

$$\mathcal{R}_z^N = \sum_{k=0}^{n-1} \sum_{B \in \mathbb{BD}_k(z)} a_B. \tag{1.1}$$

For any rectangle $V \subseteq \mathbb{Z}^2$, we denote by Π_{LR}^V the collection of all paths π in V (considered as a subgraph of $\mathbb{Z}^2$) that connects the left boundary of V to its right boundary. We refer to such a path as a (left-right) *crossing* of V. For a fixed $\gamma > 0$ we define

$$D_{\gamma,LR}(V_N) = \min_{\pi \in \Pi_{LR}^{V_N}} \sum_{z \in \pi} \mathrm{e}^{\gamma \mathcal{R}_z^N},$$

to be the first passage percolation distance between the two boundaries of V_N where we assign each vertex v a weight of $e^{\gamma \mathcal{R}_v^N}$.

Theorem 1.1. *For a fixed γ_0 that is small, and all $\gamma < \gamma_0$, we have that*

$$\mathbb{E} D_{\gamma,LR}(V_N) = O(N^{1-\gamma^2/10}).$$

*Partially supported by NSF grant DMS-1455049 and Alfred Sloan fellowship.

Remark 1.2. Our proof in fact gives an upper bound on the expected weight for the geodesic crossing through a rectangle. As a result, one can show that the expected weight for the geodesic connecting two fixed vertices has exponent strictly less than 1, by constructing a sequence of ($\log N$ many) rectangles with geometrically growing size that connect these two vertices.

1.1 A brief discussion on motivation and background

In a broad context, our work is motivated by studying first passage percolation on random media with heavy correlation. The main contribution of our work, is to demonstrate an instance that in a hierarchical random field the exponent of the FPP distance can be strictly less than 1 (despite the fact that the a straight line has weight with exponent strictly larger than 1, and that the majority of the vertices are of values $O(\sqrt{\log N})$). This is, of course, rather different from the classical FPP where the edge/vertex weights are independent and identically distributed. We refer to [2, 11] and references therein for a review of classical first passage percolation.

In the specific instance of branching random walk, our work is closely related to the first passage percolation problem when the vertex weight is given by exponentiating a two-dimensional discrete Gaussian free field (GFF). Inspired by the current work, a similar upper bound on the FPP distance in this case has been recently obtained in [6]. The FPP on two-dimensional discrete GFF, as we expect, is of fundamental importance in understanding the random metric associated with the Liouville quantum gravity (LQG) [17, 10, 18]. We remark that the random metric of LQG is a major open problem, even "just" to make rigorous sense of it. In a recent series of works of Miller and Sheffield, much understanding has been obtained (more on the continuum set up) in the special case of $\gamma = \sqrt{8/3}$; see [15, 14] and references therein. Another recent work [12] has provided some bounds on the scaling exponent for a type of LQG metric, though we emphasize that their definition is of very different flavor from the FPP perspective considered in our paper (in particular, no mathematical connection can be drawn between these works at the moment). We refrain ourselves from an extensive discussion on background of LQG in this article. As a final remark to the connection of LQG metric, we note that there are other candidate discrete metric (associated with LQG or GFF) that have been proposed, and it is possible that these metrics with suitable normalization would give a more desirable scaling limit than that of FPP. However, we feel that in the level of precision of the present article and that of [6], it is quite possible that the fundamental mathematical structures (and thus obstacles) are common for these related notions of metrics.

BRW is perhaps the simplest construction that approximates a log-correlated Gaussian field, of which the GFF is a special instance. A number of other properties, especially those regarding to the extreme values of the field, have been proved to exhibit universal behavior among log-correlated Gaussian fields (c.f., [13, 7]). Indeed, it is typical that properties were first proved for BRW, and then later for GFF; and it is also typical that the proof in the case of GFF were substantially enlightened by the understanding on BRW. However, caution is required when drawing heuristic conclusion on GFF based on BRW for FPP: as demonstrated in [8], there exists a family of log-correlated Gaussian fields whose exponent for FPP can be made arbitrarily close to 1 (as K grows to ∞) if one is allowed to perturb the covariance entry-wise by an additive amount that is at most K (and thus the scaling exponent for FPP is non-universal among log-correlated fields). That being said, we remark that while the mathematical details in [6] are substantially more involved and delicate, these two papers do share the same very basic multi-scale analysis proof framework.

1.2 Main ideas of our approach

Our approach is inspired by the rescaling argument employed in the proof of the connectivity of fractal percolation process [3, 5, 4], the idea of which went back to [1]. For the purpose of induction, we will in fact work with crossings through a rectangle rather than a square. In what follows, we should consider γ as a small positive number (but fixed as $N \to \infty$).

We wish to carry out a inductive construction and proof for light path crossing the rectangle: we will take advantage of the hierarchical structure of BRW and will construct the BRW along the way as we construct light path in different scales. In order to properly describe our procedure, we need a number of definitions. Let $\Gamma = \Gamma(\gamma) \in [\alpha/\gamma^2, \alpha/\gamma^2 + 2]$ be an odd, positive integer for some α to be selected, and let $V_N^\Gamma = ([0, \Gamma N - 1] \times [0, N - 1]) \cap \mathbb{Z}^2$. The reason for choosing Γ odd is mere technical convenience which will become apparent later. We will also work under the assumption that $\gamma \ll 1/\alpha \ll 1$ (small or large enough for our bounds or inequalities to hold although we keep this implicit in our discussions). We define a Gaussian process $\{\mathcal{R}_z^N : z \in V_N^\Gamma\}$ on V_N^Γ as follows:

- $\{\mathcal{R}_z^N : z \in V_{N;j}^\Gamma\}$ is a BRW on $V_{N;j}^\Gamma = ([(j-1)N, jN - 1] \times [0, N - 1]) \cap \mathbb{Z}^2$ where $j \in [\Gamma]$.
- $\{\mathcal{R}_z^N : z \in V_{N;j}^\Gamma\}$'s are independent of each other for $j \in [\Gamma]$.

So $\{\mathcal{R}_z^N : z \in V_N^\Gamma\}$ is basically a concatenation of Γ independent BRWs placed side by side. We can view V_N^Γ as a $\Gamma N \times N$ rectangle divided into Γ cells of dimension $N \times N$. Similarly, V_{2N}^Γ can be viewed as a $2\Gamma \times 2$ rectangle divided into 4 sub-rectangles of dimension $\Gamma N \times N$ each of which is a copy of V_N^Γ (see Figure 1 below for an illustration).



Figure 1 – $(n+1)$**-th stage of the recursive construction of BRW on** V_{2N}^Γ. Here $\Gamma = 3$.

We wish to construct a light path in V_{2N}^Γ based on constructions in V_N^Γ. As the path crosses through the rectangle of V_{2N}^Γ, we will make a choice of whether the path will stay in the upper or lower sub-rectangles based on the Gaussian values associated with the boxes of dimension $N \times N$, and thus we will switch back and forth between top and bottom layers. Our goal is to show that the expected weight of the crossing we constructed expands by a factor less than 2 when the size of the rectangles doubles. Apart from doubling of dimension at each level,there are a number of reasons that the weight will expand: (1) For standard Gaussian variable, we have $\mathbb{E}e^{\gamma X} = 1 + \frac{\gamma^2}{2} + O(\gamma^3)$;

(2) We will need to make a construction to connect the paths on the left half and the right half of the rectangle; (3) Every time we switch between top and bottom layer, we will have to add a top-down crossing in the switching location. What we would possibly gain is from the "variation" of these Gaussian variables associated with $N \times N$ boxes, and thus choosing the favorable layer would reduce the weight of our path. There are a couple of subtle points in carrying out this plan, as we explain now.

In order to see how switching will reduce our cost, we first consider a toy problem where we simply compute $\mathbb{E}\min(X,Y)$ for independent standard Gaussian variables X and Y. It turns out for this purpose it is more convenient to write $X = \frac{X+Y}{2} + \frac{X-Y}{2}$ and $Y = \frac{X+Y}{2} - \frac{X-Y}{2}$, and thus $\mathbb{E}\min(X,Y) = -\mathbb{E}|\frac{X-Y}{2}| = \frac{-1}{\sqrt{\pi}}$. If we carry out inductive procedure in this fashion, what we gain from the switching will not be able to beat the aforementioned expansions. This is because, in the language of the toy model, $\min(X,Y)$ is itself a random variable that can be written as $\frac{X+Y}{2} - \frac{|X-Y|}{2}$, where we gain $-\frac{1}{\sqrt{\pi}}$ by averaging over $(X-Y)$ but we get nothing by averaging over $X+Y$ (which is independent of $(X-Y)$). In fact, the better strategy would be to save the variation from $(X+Y)$ and use it in the optimizations in the next scale. But it is inconvenient to carry out this strategy, as it would require to do an induction based on quantities beyond the expected weight of the crossings in different scales. We resolve this issue by represent the BRW in a different way, which is tailored for our application here.

Another point is that, in the aforementioned optimization procedure, one has to be rather strategic in switchings as there is a cost for that. Eventually, this is reduced to a question of computing the following *regularized total variation* for the Brownian motion, defined by

$$\mathbb{E}\max_k \max_{0=t_0<t_1<t_2<\ldots<t_k<t_{k+1}=1} \sum_{i=1}^{k+1} |B_{t_i} - B_{t_{i-1}}| - \lambda k\,. \tag{1.2}$$

Here $B_{[0,1]}$ is a standard Brownian motion, and $\lambda > 0$ is the term that measures the switching cost. We will use a recent result of [9] on the asymptotic value of (1.2) as $\lambda \to 0$.

Notation convention. In the rest of the paper we refer to the vertices of $\mathbb{Z}^2$ as *points*. For any point $z \in \mathbb{Z}^2$, the horizontal and vertical coordinates of z are denoted by z_x and z_y respectively. If $z_y = 0$, we represent z simply as z_x, which should be clear from the context. A (finite) *path* π is a finite sequence $(v_0, v_1, \cdots, v_m)$ of points such that for each $i \geq 0$, v_i and v_{i+1} are neighbors in $\mathbb{Z}^2$. In each of the pairs of non-negative integers $(n,N), (\ell,L), (\ell',L')$ and (ℓ'',L''), the second element is always assumed to be 2 raised to the power of first element. For functions $F(.)$ and $G(.)$, we write $F = O(G)$ if there exists an absolute constant $C > 0$ such that $F \leq CG$ everywhere in the domain. We write $F = \Omega(G)$ if $G = O(F)$.

Acknowledgement. We warmly thank Steve Lalley for numerous helpful discussions and constant encouragement on the project.

2 Preliminaries

2.1 Coarsening of paths and $\mathcal{L}$-segments

The *L-coarsening* of a (finite) path $\pi = (v_0, v_1, \cdots, v_m)$ in $\mathbb{Z}^2$ is defined as follows. Cover the lattice $\mathbb{Z}^2$ with $L \times L$ (non-overlapping) boxes such that one such box is V_L. Denote the center of a box B as $c(B)$. For $v \in \mathbb{Z}^2$, let $B_L(v)$ be the unique box containing v. Now define a sequence of integers $m_0, m_1, \cdots$ recursively as $m_0 = 0$ and,

$$m_j = \inf\{m \geq i \geq m_{j-1} : v_i \notin B_L(v_{m_{j-1}})\} \text{ for } j \geq 1\,.$$

Here we adopt the standard convention that infimum of an empty set is ∞. Let $p_{L,\pi}$ be the last integer j such that m_j is finite. The L-coarsening of π, denoted as $\pi_{L,coarse}$, can now be defined as the sequence $\big(c(B_L(v_{m_0})), c(B_L(v_{m_1})), \cdots, c(B_L(v_{m_{p_{L,\pi}}}))\big)$. We can also treat $\pi_{L,coarse}$ as a path in $\mathbb{R}^2$ that connects successive points in this list by a line segment. See Figure 2 for an illustration. Notice that the coarsening of a simple (i.e. self-avoiding) path is not necessarily simple. However if the L-coarsening of a (left-right) crossing π^* of V^{Γ}_{2L} is simple, then the $2L$-coarsening of π^* is also simple. We will use this fact when we carry out the induction step.
An *$\mathcal{L}$-segment at level ℓ* is a path π whose 2^ℓ-coarsening is one of two shapes shown in Figure 3.



Figure 2 – **L-coarsening of a path**. The dashed path on the right represents the L-coarsening of the solid path on the left.



Figure 3 – **Two possible 2^ℓ-coarsenings of an $\mathcal{L}$-segment at level ℓ.**

2.2 Switched sign construction for branching random walk

We first introduce a few more notations. As described in the introduction, we view V^{Γ}_{2N} as a $2\Gamma \times 2$ rectangle divided into 4 sub-rectangles of dimension $\Gamma N \times N$ each of which is a copy of V^{Γ}_N (see Figure 1 for an illustration). Denote these 4 subrectangles (or the copies of V^{Γ}_N) as $\{V^{\Gamma}_{N;i,k}\}_{i\in[2],k\in[2]}$ in the usual order. We can also shift the origin by $u = (\Gamma i 2^{n+1}, j2^{n+1})$ for $i, j \in \mathbb{N} \cup \{0\}$ and denote the corresponding subrectangles (respectively rectangles) by $\{V^{\Gamma,u}_{N;i,k}\}_{i\in[2],k\in[2]}$ (respectively $V^{\Gamma,u}_{2N}$). Similarly we can define the fields $\{\mathcal{R}^{N,u}_{z;i,k} : z \in V^{\Gamma,u}_{N;i,k}\}$ or $\{\mathcal{R}^{N,u}_{z} : z \in V^{\Gamma,u}_{N}\}$. The j-th constituent cell in $V^{\Gamma,u}_{N;i,k}$ (which is box of side length N) will be denoted as $B^u_{n;i,k,j}$ where $j \in [\Gamma]$.

We observe that the Gaussian field $\{\mathcal{R}_z^{2N,u} : z \in V_{2N}^{\Gamma,u}\}$ is obtained by adding $a_{B^u_{n;i,k,j}}$ to the field $\{\mathcal{R}_{z;i,k}^{N,u} : z \in V_{N;i,k}^{\Gamma,u}\}$ in the box $B^u_{n;i,k,j}$ for $i \in [2], k \in [2]$ and $j \in [\Gamma]$ (See Figure 1). We will omit the additional superscript u in all these notations when $u = 0$.

We will represent BRW by a construction in the fashion of switching-signs, which is tailored to our inductive construction for light crossings. To this end, we denote the collection of all rectangles of the form $([0, 2^\ell - 1] \times [0, 2^{\ell+1} - 1]) \cap \mathbb{Z}^2 + (i2^\ell, j2^\ell)$ by $\mathbb{BD}_{k,2}$ where i, j, ℓ are nonnegative integers. Let $A_{\ell,\Gamma}$ be the collection of all points of the form $(\Gamma i 2^\ell, j2^\ell)$. Denote by $R^u_{\ell;k,j}$ the rectangle formed by the boxes $B^u_{\ell;1,k,j}$ and $B^u_{\ell;2,k,j}$ when $x \in A_{\ell+1,\Gamma}$. Note that $R^u_{\ell;k,j} \in \mathbb{BD}_{\ell,2}$. Let $\{a_B\}_{B\in\mathbb{BD}_{k,2},k\geq 0}$ be a collection of i.i.d. standard normal random variables, that is independent of $\{a_B\}_{k\geq 0, B\in\mathbb{BD}_k}$. We will now construct a family of Gaussian fields $\{\chi_z^{L,u} : z \in V_{2^\ell}^{\Gamma,u}\}$ for $u \in A_{\ell,\Gamma}$ and $\ell = 0, 1, 2, \ldots, n$ recursively as follows:

- $\chi_z^{1,u} = 0$ for all $z \in V_1^{\Gamma,u}$.
- Let $u \in A_{\ell+1,\Gamma}$ and $\{\chi_{z;i,k}^{L,u} : z \in V_{L;i,k}^{\Gamma,u}\}$ be the field on $V_{L;i,k}^{\Gamma,u}$ constructed in level ℓ. For $z \in B^u_{\ell;2,k,j}$, define

$$\chi_z^{2L,u} = \chi_{z;i,k}^{L,u} + (-1)^{(k-1)\Gamma+j} b_\ell a_{\tilde{R}^u_{\ell;k,j}} + (-1)^i c_\ell a_{B^u_{\ell;2,k,j}} ,$$

  where $b_\ell = \sqrt{(1 - 4^{-\ell})/3}, c_\ell = \sqrt{2(1 - 4^{-\ell})/3}$ and $\tilde{R}^u_{\ell;k,j}$ is the rectangle defined as follows:

$$\tilde{R}^u_{\ell;k,j} = \begin{cases} R^u_{\ell;k,j} & \text{if } (k-1)\Gamma + j \text{ is odd}, \\ R^u_{\ell;k,j-1} & \text{if } (k-1)\Gamma + j \text{ is even and } j > 1, \\ R^u_{\ell;k-1,\Gamma} & \text{if } (k-1)\Gamma + j \text{ is even and } j = 1. \end{cases}$$

For future reference we will denote $Z_{\ell;i,k,j} = (-1)^{(k-1)\Gamma+j} b_\ell a_{\tilde{R}_{\ell;k,j}} + (-1)^i c_\ell a_{B_{\ell;2,k,j}}$. Similarly as before, we will drop the superscript u whenever $u = 0$. See Figure 4 for an illustration of this construction at level $\ell + 1$. Notice that since $c_\ell^2 = 2b_\ell^2$, distribution of the field $\{\chi_z^N : z \in V_{N;j}^\Gamma\}$ is symmetric with respect to symmetries of the box $V_{N;j}^\Gamma$. This fact will be used later when we construct crossings through $V_{N;j}^\Gamma$'s in vertical direction. Finally, we define

$$\tilde{\chi}_z^{N,u} = b_n a_{V^u_{N;j}} + \chi_z^{N,u} \text{ for } u \in A_{n,\Gamma} \text{ and } z \in V_{N;j}^{\Gamma,u} = u + V_{N;j}^\Gamma .$$

Lemma 2.1. *The Gaussian fields $\{\tilde{\chi}_z^N : z \in V_N^\Gamma\}$ and $\{\mathcal{R}_z^N : z \in V_N^\Gamma\}$ are identically distributed.*

Proof. It is not difficult to check that the fields $\{\tilde{\chi}_z^N : z \in V_{N;j}\}$'s are independent and identically distributed for $j \in [\Gamma]$. Hence it suffices to verify that we have the correct covariances between field values at all pairs of vertices inside V_N. Toward this end take a pair (u, v) of vertices in V_N which were separated until level ℓ, i.e., $\mathrm{Cov}(\mathcal{R}_u^N, \mathcal{R}_v^N) = n - \ell$. The covariance between χ_u^N and χ_v^N is given by

$$\mathrm{Cov}(\chi_u^N, \chi_v^N) = -(1 - 4^{-\ell})/3 + \textstyle\sum_{\ell+1\leq\ell\leq n}(1 - 4^{-\ell}) = n - \ell - (1 - 4^{-n})/3 .$$

Consequently $\mathrm{Cov}(\tilde{\chi}_u^N, \tilde{\chi}_v^N) = \mathrm{Cov}(\chi_u^N, \chi_v^N) + (1 - 4^{-n})/3 = n - \ell$. Hence the lemma follows. □

Remark 2.2. It is clear that it suffices to work with $\{\chi_z^N : z \in V_N^\Gamma\}$ rather than the BRW in what follows. Using this construction, in every scale of our optimization, we crucially have effective variance $c_\ell^2 \approx \frac{2}{3}$ that is larger than $1/2$ (which would be the variance we effectively used if we work with the more canonical construction of BRW).


$V^{\Gamma}_{L;1,1}$
$-b_\ell a_{R_{\ell;1,1}} - c_\ell a_{B_{\ell;2,1,1}}$
$b_\ell a_{R_{\ell;1,1}} - c_\ell a_{B_{\ell;2,1,2}}$
$-b_\ell a_{R_{\ell;1,3}} - c_\ell a_{B_{\ell;2,1,3}}$
$-b_\ell a_{R_{\ell;1,1}} + c_\ell a_{B_{\ell;2,1,1}}$
$b_\ell a_{R_{\ell;1,1}} + c_\ell a_{B_{\ell;2,1,2}}$
$-b_\ell a_{R_{\ell;1,3}} + c_\ell a_{B_{\ell;2,1,3}}$
$V^{\Gamma}_{L;2,1}$

Figure 4 – **Construction of the field χ_z^{2L} on $V^{\Gamma}_{L;i,1}$.** Here $\Gamma = 3$. The red lines indicate the boundary of $R_{\ell;1,1}$.

2.3 Regularized total variation of Brownian motion

In this subsection we present a result from [9] that provides a lower bound on the functional defined in (1.2). This is only one part of the main result in [9] which also provides an upper bound.

Theorem 2.3. *Let $\{B_t\}_{0\le t\le 1}$ be a Standard Brownian Motion and $1 > \lambda > 0$. For any partition $P = [t_0, t_1, \cdots, t_{k+1}]$ of $[0,1]$ where $0 = t_0 < t_1 < \cdots < t_{k+1} = 1$, define the quantity $\Phi_{P,\lambda}$ as*

$$\Phi_{P,\lambda} = \textstyle\sum_{0\le i\le k} |B_{t_{i+1}} - B_{t_i}| - \lambda k\,.$$

Then there exists a (random) partition $\mathcal{P} = [t_0^*, t_1^*, \cdots, t_{k^*+1}^*]$ of $[0,1]$ such that the following conditions hold:

(a) The random vector $(t_1^*, \cdots, t_{k^*}^*, k^*)$ is measurable relative to $\sigma(\{B_t\}_{0\le t\le 1})$.

(b) $\mathbb{E}\Phi_{P,\lambda} \ge \frac{1}{\lambda}$.

(c) $B_{t_1^*} - B_{t_0^*}, B_{t_2^*} - B_{t_1^*}, \cdots, B_{t^*_{k^*+1}} - B_{t_{k^*}^*}$ have alternating signs. Furthermore $\mathbb{P}(B_{t_1^*} - B_{t_0^*} > 0) = 0.5$.

(d) $|B_{t_{j+1}^*} - B_{t_j^*}| \ge \lambda$ for all $1 \le j \le k^* - 1$.

3 Inductive construction for light crossings

3.1 The induction hypotheses

We will construct a light (left-right) crossing through V_L^Γ (or equivalently $V_L^{\Gamma,u}$) inductively for $L = 2^\ell$ and $\ell = 0, \dots, n$. Throughout this section whenever we refer to a crossing of $V_L^{\Gamma,u}$, it is implicitly assumed that the underlying field is $\{\chi_z^{L,u} : z \in V_L^{\Gamma,u}\}$ unless stated otherwise. For technical convenience, let us assume that for each $\ell \geq 0$ and $u \in A_{\ell+1,\Gamma}$ we have an independent standard Brownian motion $\{W_t^{u,\ell}\}_{0\leq t\leq 2\Gamma}$ that is also independent with $\{a_B\}_{B\in\mathbb{BD}_{k,2},k\geq 0}$. We sample $a_{B^u_{\ell;2,k,j}}$'s as increments of this process at appropriate time points. We are now ready to state our induction hypotheses. For $\ell = 0$, we have only one option, i.e., to take the straight line as our crossing. This crossing has weight exactly Γ. Suppose that for each $1 \leq \ell' \leq \ell$ and $u \in A_{\ell',\Gamma}$, we have a crossing $\pi^{*,u,\ell'}$ of $V_{2^{\ell'}}^{\Gamma,u}$, identically distributed for different u, that satisfies the following properties:

1. $\pi^{*,u,\ell'}_{L'/2,coarse}$ is simple and measurable relative to $\sigma\big(\{W_t^{u,\ell'-1}\}_{0\leq t\leq 2\Gamma}\big)$.
2. Law of $\pi^{*,u,\ell'}_{L'/2,coarse}$ is invariant with respect to reflection about the line $y = u_y + 2^{\ell'-1} - 0.5$ i.e. the horizontal line passing through the center of $V_{2^{\ell'}}^{\Gamma,u}$.
3. For $j \in [\Gamma]$, denote the line $x = u_x + jL' - 1$ as $\mathbb{L}_{jL'}$. Let $z^u_{j,\ell'}$ be the point in the set $\{u, u+(0,L'/2), u+(\Gamma L'/2, 0), u+(\Gamma L'/2, L'/2)\}$ such that the point where $\pi^{*,u,\ell'}$ hits $\mathbb{L}_{jL'}$ for the last time lies within $V^{\Gamma,z^u_{j,\ell'}}_{L'/2}$. Also for $j \in \mathbb{N}^+$, let $v^u_{j,L',L''}$ be the last point along $\pi^{*,u,\ell'}_{L'',coarse}$ that lies to the left of $\mathbb{L}_{jL'}$. Then $v^u_{j,L',L'_{100}} = v^{z^u_{j,\ell'}}_{2j,L'/2,L'_{100}}$, i.e., the crossings $\pi^{*,u,\ell'}$ and $\pi^{*,z^u_{j,\ell'},\ell'-1}$ hit $\mathbb{L}_{jL'}$ for the last time in the same $L'_{100}\times L'_{100}$ box. Here $L'_{100} = \max(2^{-100}L', 1)$.
4. $\mathbb{E}(\sum_{z\in\pi^{*,u,\ell'}} \mathrm{e}^{\gamma\chi_z^{\ell'}}) \leq 2\Gamma(2 - 0.2\gamma^2)^{\ell'}$.

One of the consequences of our induction hypotheses, which is crucial for our analysis, is that $v^u_{j,L,\delta L}$ is uniformly distributed for all $\ell' \leq \ell$ as given by the following lemma.

Lemma 3.1. *Let $R^u_{\ell;i,k,j}$ be the unique $L_{100} \times L$ rectangle that shares its right boundary with with $B^u_{\ell;i,k,j}$. Also let $\{B^u_{\ell;i,k,j,m}\}_{1\leq m\leq L/L_{100}}$ be the collection of boxes in $\mathbb{BD}_{\log_2 L_{100}}$ that comprise $R^u_{\ell;i,k,j}$. Now if the induction hypotheses 1, 2 and 3 hold for all $\ell' \leq \ell$, then $v^u_{j,L,L_{100}}$ is distributed uniformly on the set $\{c^u_{\ell;i,k,j,m}\}_{1\leq m\leq L/L_{100}}$, where $c^u_{\ell;i,k,j,m}$ is the center of the box $B^u_{\ell;i,k,j,m}$ (see Figure 5).*

Proof. We get, from hypothesis 2, that for any $2 \leq L' \leq L, j' \in [\Gamma]$ and $x \in A_{\ell',\Gamma}$, $v^u_{j',L',L'/2}$ is equally likely to be above or below the line $y = u_y + 2^{\ell'-1} - 0.5$. We can recursively apply this argument for all the levels down to $\log_2 L_{100}$ in view of hypothesis 1 which gives that the $L'/2$-coarsenings of crossings at level ℓ' are independent for different ℓ'. Combining these observations with hypothesis 3 completes the proof of the lemma. □

3.2 The induction step

Now we will carry out the induction step. It suffices to produce a crossing for $u = 0$. As usual, we will drop u from all the superscripts when u is origin. Before going to more detailed discussions let us give a broad overview of the strategy that we follow.



Figure 5 – **The set of points** $\{c_{\ell;i,k,j,m}\}_{1\leq m\leq 2^{100}}$.

First we decide the L-coarsening of $\pi^{*,\ell+1}$. It would consist of a sequence $H_1, H_2, \cdots, H_{R+1}$ of horizontal segments (possibly of length 0) from left to right such that every two successive segments are connected by a vertical segment. See Figure 6 for an illustration. Notice that this is the only possible option for $\pi^{*,\ell+1}_{L,coarse}$, if it has to be simple as required by induction hypothesis 1. So we can "encode" $\pi^{*,\ell+1}_{L,coarse}$ as a sequence of $\{1,2\}$ valued random variables $\{i_{j,k}\}_{j\in[\Gamma],k\in[2]}$, where $i_{k,j}=1$ or 2 accordingly as $\pi^{*,\ell+1}_{L,coarse}$ enters the rectangle $R_{\ell;k,j}$ through $B_{\ell;1,k,j}$ or $B_{\ell;2,k,j}$ respectively. We refer to this as a *switching strategy*. Having defined L-coarsening of the crossing, we now move on to actual construction. From the previous level we have four crossings $\{\pi^{*,\ell}_{i,k}\}_{i\in[2],k\in[2]}$ respectively in $\{V^{\Gamma}_{L;i,k}\}_{i\in[2],k\in[2]}$. We "join" the paths $\pi^{*,\ell}_{i,1}$ and $\pi^{*,\ell}_{i,2}$ into a crossing $\pi^{*,\ell+1}_{i}$ of V^{Γ}_{2L} for $i\in[2]$. Since L-coarsening of crossings at level ℓ are also simple, it follows that the vertices of $\pi^{*,\ell+1}_{i}$ that lie within some $B_{\ell;i,k,j}$ define a subpath of $\pi^{*,\ell+1}_{i}$. Now we replace each H_r with the subpath $h^{*,\ell+1}_{r}$ of $\pi^{*,\ell+1}_{i}$ that is contained within the $L\times L$ boxes that intersect H_r. Here i is 1 or 2 accordingly as H_r lies in the top or bottom row of V^{Γ}_{2L} (See Figure 6 and the discussions preceding Figure 1). Finally we connect $h^{*,\ell+1}_{r}$ and $h^{*,\ell+1}_{r+1}$ by an appropriate $\mathcal{L}$-segment $v^{*,\ell+1}_{r}$ at level ℓ. The paths $h^{*,\ell+1}_{1}, v^{*,\ell+1}_{1}, h^{*,\ell+1}_{2}, \cdots, v^{*,\ell+1}_{R}, h^{*,\ell+1}_{R+1}$ define the crossing $\pi^{*,\ell+1}$. Note that L-coarsening of $\pi^{*,\ell+1}$ is the very one that we started with.

We also need some notations in order to track the change in expected weight of crossings between



Figure 6 – **L-coarsening of $\pi^{*,\ell+1}$.** Here $\Gamma = 3$ and $R = 2$.

two levels. To this end define,

$$D_{\gamma,\ell} = \sum_{v\in\pi^{*,\ell}} e^{\gamma\chi_v^L}\,,\ \text{and}\ d_{\gamma,\ell} = \mathbb{E}D_{\gamma,\ell}$$
$$D_{\gamma,\ell,j} = \sum_{v\in\pi^{*,\ell}\cap V_{L,j}} e^{\gamma\chi_v^L},\ \text{and}\ d_{\gamma,\ell,j} = \mathbb{E}D_{\gamma,\ell,j}\ \text{for}\ j\in[\Gamma]\,.$$

Let $\tilde{D}_{\gamma,\ell;k}$ the weight contributed to $\pi^{*,\ell+1}$ by the paths $\pi_{i,k}^{*,\ell}$'s. From what we described in the previous paragraph, we may bound $\mathbb{E}\tilde{D}_{\gamma,\ell;k}$ as

$$\mathbb{E}\tilde{D}_{\gamma,\ell;k} \le \mathbb{E}\sum_{j=1}^{\Gamma} d_{\gamma,\ell,j}\big(1+\gamma Z_{\ell;i_{j,k},k,j} + \frac{a_{\ell+1}^2\gamma^2}{2}\big) + \mathbb{E}\mathrm{Err}_{\gamma,\ell;k,1} + \mathbb{E}\mathrm{Err}_{\gamma,\ell;k,2}\,, \tag{3.1}$$

where $a_{\ell+1}^2 = b_{\ell+1}^2 + c_{\ell+1}^2 = 1 - 4^{-(\ell+1)}$, $\mathrm{Err}_{\gamma,\ell;k,1} = \frac{\gamma^2}{2}\sum_{j=1}^{\Gamma} d_{\gamma,\ell,j}(Z_{\tilde{\ell};i_{j,k},k,j}^2 - a_{\ell+1}^2)$ and

$$\mathrm{Err}_{\gamma,\ell;k,2} = \sum_{j=1}^{\Gamma} d_{\gamma,\ell,j}\big(\mathrm{e}^{\gamma Z_{\ell;i_{j,k},k,j}} - (1+\gamma Z_{\ell;i_{j,k},k,j} + \frac{Z_{\ell;i_{j,k},k,j}^2\gamma^2}{2})\big)\,.$$

It is easy to see that $\mathbb{E}|\mathrm{Err}_{\gamma,\ell;k,2}| = d_{\gamma,\ell}O(\gamma^3)$. On the other hand, we get the following from expanding the terms in $\mathrm{Err}_{\gamma,\ell;k,1}$:

$$\mathrm{Err}_{\gamma,\ell;k,1} = \frac{b_\ell^2\gamma^2}{2}\sum_{j=1}^{\Gamma} d_{\gamma,\ell,j}(a_{\tilde{R}_{\ell;k,j}}^2 - 1) + \frac{c_\ell^2\gamma^2}{2}\sum_{j=1}^{\Gamma} d_{\gamma,\ell,j}(a_{B_{\ell;2,k,j}}^2 - 1)$$
$$+\gamma^2 b_\ell c_\ell(-1)^{(k-1)\Gamma}\sum_{j=1}^{\Gamma} d_{\gamma,\ell,j}a_{\tilde{R}_{\ell;k,j}}(-1)^{j+i_{j,k}}a_{B_{\ell;2,k,j}}\,.$$

Expectations of the first two terms in the expansion above are obviously 0. As to the third term, recall that our switching mechanism is independent of the random variables $a_{\tilde{R}_{\ell;k,j}}$'s by induction hypothesis 2. Consequently $\mathbb{E}\mathrm{Err}_{\gamma,\ell;k,1} = 0$. Thus we only need to reckon with the first term in the right hand side of (3.1) so far as the contributions from $\pi_{i,k}^{*,\ell}$'s are concerned. This will, of course, depend on the particular switching strategy we adopt. Finally we need to add the contributions from $v_r^{*,\ell+1}$'s and the extra crossings that we make in order to link $\pi_{i,1}^{*,\ell}$ and $\pi_{i,2}^{*,\ell}$ for $i\in[2]$.

Now we carry out the detailed computations. It is expected that $d_{\gamma,\ell,j}$ is approximately uniform over $j \in [\Gamma]$. However, proving this requires some effort. It turns out easier to treat the "hypothetical" case that $d_{\gamma,\ell}$ is dominated by a small number of $d_{\gamma,\ell,j}$'s. We also include a case to deal with small values of ℓ when c_ℓ^2 is not big enough for our "main" strategy to work.

Case 1. $\sum_{j=1}^{\Gamma} d_{\gamma,\ell,j}\mathbf{1}_{\{d_{\gamma,\ell,j}\geq \Gamma^{-2/3}d_{\gamma,\ell}\}} \geq d_{\gamma,\ell}\Gamma^{-1/10}$. It is easy to see that

$$\sum_{j=1}^{\Gamma} d_{\gamma,\ell,j}^2 \geq d_{\gamma,\ell}^2\Gamma^{-13/15}\,. \tag{3.2}$$

In this case our switching strategy is simple. For each $k = 1, 2$, we choose the rectangle that minimizes $\sum_{j=1}^{\Gamma} d_{\gamma,\ell,j}Z_{\ell;i,k,j}$. Thus, we have for $k \in [2]$

$$\begin{aligned}\mathbb{E}\tilde{D}_{\gamma,\ell;k} &\leq d_{\gamma,\ell}(1+\gamma^2/2+O(\gamma^3)) + \mathbb{E}\min_{i=1,2}\sum_{j=1}^{\Gamma} d_{\gamma,\ell,j}\gamma Z_{\ell;i,k,j}\\ &\leq d_{\gamma,\ell}(1+O(\gamma^2)-\Omega(\gamma)\Gamma^{-13/30}) \leq d_{\gamma,\ell}(1-\Omega(\gamma^{1.9}))\,,\end{aligned}$$

where the last inequality follows from our assumption that $\Gamma \geq \alpha/\gamma^2$. Let $u_k \in A_{\ell,\Gamma}$ be such that V_L^{Γ,u_k} is the minimizer. In order to connect $\pi^{*,u_1,\ell}$ and $\pi^{*,u_2,\ell}$ into a crossing of V_{2L}^{Γ}, we need some additional paths. To this end we add two vertical crossings through the rectangles $V_{\text{left}} = \big([(\Gamma-1)L, \Gamma L-1]\times[0,2L-1]\big)\cap\mathbb{Z}^2$ and $V_{\text{right}} = \big([\Gamma L, (\Gamma+1)L-1]\times[0,2L-1]\big)\cap\mathbb{Z}^2$. We also add two horizontal crossings through the rectangles $V_{\text{up}} = \big([(\Gamma-1)L, (\Gamma+1)L-1]\times[L,2L-1]\big)\cap\mathbb{Z}^2$ and $V_{\text{down}} = \big([(\Gamma-1)L, (\Gamma+1)L-1]\times[0,L-1]\big)\cap\mathbb{Z}^2$. All these crossings have minimum weights. We can now form $\pi^{*,\ell+1}$ by concatenating $\pi^{*,u_1,\ell}, \pi^{*,u_2,\ell}$ and these four crossings. See Figure 7 for an illustration. It is obvious from the construction that $\pi^{*,\ell+1}_{L,coarse}$ is simple, symmetric in law with respect to the line $y = L-0.5$ and entirely determined by the random variables $\{a_{B_{\ell;2,k,j}}\}_{k\in[2],j\in[\Gamma]}$. The only possible case where induction hypothesis 3 could fail would be the line $\mathbb{L}_{\Gamma L}$. But we do not need to consider this line as Γ is odd. Finally notice that, since $\Gamma \gg 1$ and minimum crossing weights through adjacent rectangles are super-additive, each of the four additional crossings will have expected weight bounded by, say, $\frac{2.1}{\Gamma}d_{\gamma,\ell}$. Using the fact that $\alpha \gg 1$, we then deduce that in this case

$$d_{\gamma,\ell+1} \leq (2-\gamma^2/2)d_{\gamma,\ell} \leq 2\Gamma(2-0.2)^{\ell+1}\,.$$



Figure 7 – **The construction of $\pi^{*,\ell+1}$ for Case 1.** Here $\Gamma = 3$. The segments of $\pi^{*,\ell+1}$ have been colored in red.

Case 2. If $\ell < 60$, we use the strategy described in Case 1. It then follows from the previous discussions that

$$d_{\gamma,\ell+1} \leq (2+2\gamma^2)d_{\gamma,\ell}\,.$$

Since $d_{\gamma,0} = \Gamma$, we get by repeated application of the previous inequality

$$d_{\gamma,\ell+1} \le \Gamma(2+2\gamma^2)^{\ell+1} \le 2\Gamma(2-0.2\gamma^2)^{\ell+1}\,.$$

Case 3. $\sum_{j=1}^{\Gamma} d_{\gamma,\ell,j}\mathbf{1}_{d_{\gamma,\ell,j}\ge\Gamma^{-2/3}d_{\gamma,\ell}} \le d_{\gamma,\ell,j}\Gamma^{-1/10}$ and $\ell \ge 60$. This is the main and real case. We will also assume that

$$d_{\gamma,\ell} \ge 2\Gamma(2-0.2\gamma^2)^{\ell}(1-\gamma^2)\,,$$

for otherwise we can simply follow the strategy in Case 1. In this case our switching strategy will in fact be informed by the way we plan to contruct the $\mathcal{L}$-segments $v_r^{*,\ell+1}$'s. So we first give a detailed description of the latter. In what follows, we may assume without any loss of generality that $\delta L \ge 2$ for $\delta = 2^{-100}$. We refer the reader to Fig 8 for an illustration of the construction.
For each $r \le R$, let $V_{L;U(r)}$ and $V_{L;D(r)}$ respectively denote the $L\times L$ boxes in the top and bottom row of V_{2L}^{Γ} containing the endpoints of H_r and H_{r+1}. Call the $L\times 2L$ rectangle formed by $V_{L;U(r)}$ and $V_{L;D(r)}$ as $V_{L;r}$. Suppose the right boundary of $V_{L;r}$ lies along the line $\mathbb{L}_{j_rL}$ where $j_r \in [2\Gamma-1]$. It suffices to consider the case when $j_r \le \Gamma$. Let $v^u_{j_r,L,\delta L}$ and $v^d_{j_r,L,\delta L}$ respectively denote the last points along $\pi^{*,L,\ell}_{\delta L,coarse}$ and $\pi^{*,\ell}_{\delta L,coarse}$ that lie to the left of $\mathbb{L}_{j_rL}$. Then $v^u_{j_r,L,\delta L}$ and $v^d_{j_r,L,\delta L}$ are centers of two $\delta L\times\delta L$ boxes, say, $A^u_{j_r,L}$ and $A^d_{j_r,L}$ respectively. Let $\tilde{A}^u_{j_r,L}$ (respectively $\tilde{A}^d_{j_r,L}$) be the box around $A^u_{j_r,L}$ (respectively $\tilde{A}^d_{j_r,L}$) of side length $2\delta L$. Then we construct light contours in each of the annuli $\tilde{A}^u_{j_r,L}\setminus A^u_{j_r,L}$ and $\tilde{A}^d_{j_r,L}\setminus A^d_{j_r,L}$. These can easily be achieved by concatenating four minimum weight crossings in four rectangles. In addition, we let $\tilde{A}_r$ be a rectangle whose left top corner is $v^u_{j_r,L,\delta L}+(0.5,\delta L-0.5)$ and right bottom corner is $v^d_{j_r,L,\delta L}+(0.5,-\delta L+0.5)$. We construct a minimum weight vertical crossing through $\tilde{A}_r$. Finally we concatenate these light crossings into $v_r^{*,\ell+1}$. The total expected weight of these crossings can be bounded by our induction hypotheses. Notice that we can always concatenate these paths in a way so that our construction of $v_r^{*,\ell+1}$ obeys induction hypothesis 3.

Let us now try to bound the expected weight of $v_r^{*,\ell+1}$ where we assign vertex weight based on variables *associated with boxes of side length* $\le L$. Notice that this does not require the knowledge of particular switching strategy we employ. Denote by D_r the total weight of the light crossings in the two annuli (where we assign vertex weight based variables associated with boxes of side length $\le \delta L/2$). Since the δL coarsening of $\pi^{*,u,\ell'}$ is independent of all the Gaussians at level lower than $\delta\ell$, we see that

$$\mathbb{E}D_r \le \frac{33 d_{\gamma,\ell-\log_2(2/\delta)}}{\Gamma}\,. \tag{3.3}$$

Let $\tilde{D}_r$ denote the weight of the vertical crossing through $\tilde{A}_r$ (where again, we assign the vertex weight based on Gaussian variables associated with boxes of side length $\le \delta L/2$). Clearly we have

$$\mathbb{E}\tilde{D}_r \le \frac{\big(3\delta L + \mathbb{E}|v^u_{j_r,L,\delta L} - v^d_{j_r,L,\delta L}|\big)}{(\delta L/2)\Gamma} d_{\gamma,\ell-\log_2(2/\delta)}\,. \tag{3.4}$$

In both (3.3) and (3.4), we have silently used the fact that expected minimum weights of crossings through $V^m_{L'}$'s form a superadditive sequence in m. The following lemma says that we do not lose much in terms of the expected costs of these crossings after we take into account the contributions from Gaussians appearing in higher levels.

Lemma 3.2. *Let $X_1,\dots,X_m$ i.i.d. standard Gaussian variables, where $m \le (1/\gamma)^{100}$ and U be a nonnegative random variable bounded by C. Then*

$$\mathbb{E}(U\max_i \mathrm{e}^{\gamma X_i}) \le \mathbb{E}U\big(1+O(\gamma\log(1/\gamma))\big) + C\mathrm{e}^{-\Omega(\log(1/\gamma))}\,.$$



Figure 8 – **The construction of $v_r^{*,\ell+1}$.** The two big boxes on the left define the rectangle $V_{L;r}$, the top and bottom one being $V_{L;U(r)}$ and $V_{L;D(r)}$ respectively. The two boxes filled with light blue are $A^u_{j_r,L}$ (top) and $A^d_{j_r,L}$ (bottom). The annuli $\tilde{A}^u_{j_r,L} \setminus A^u_{j_r,L}$ and $\tilde{A}^d_{j_r,L} \setminus A^d_{j_r,L}$ (or portions thereof) are indicated by light brown. The path colored in red is $v_r^{*,\ell+1}$. The green paths represent $h_r^{*,\ell+1}$ and $h_{r+1}^{*,\ell+1}$.

Proof. Let $M = \max_i X_i$. It is clear that $\mathbb{E}M \le \sqrt{200\log(1/\gamma)}$, and also that $\mathbb{P}(M \ge \mathbb{E}M + u) \le e^{-u}$. The conclusion follows by a straightforward computation. □

Combining (3.3), (3.4), Lemmas 3.1 and 3.2, we get that the expected weight for each "switching gadget"is bounded by $(1+10\delta)(2-0.2\gamma^2)^\ell \le \frac{1+20\delta}{\Gamma} d_{\gamma,\ell}$, where we use the fact that $\gamma \ll 1$. Recall the expression for $\mathbb{E}\tilde{D}_{\gamma,\ell;k}$ as given in (3.1) and the fact that

$$Z_{\ell;i,k,j} = (-1)^{(k-1)\Gamma+j} b_\ell a_{\tilde{R}_{\ell;k,j}} + (-1)^i c_\ell a_{B_{\ell;2,k,j}} .$$

It is then natural to consider switching strategies $\{i_{j,k}\}_{j\in[\Gamma],k\in[2]}$ that gives a small value of the following quantity:

$$c_\ell \gamma \sum_{j=1}^{\Gamma} d_{\gamma,\ell,j} (-1)^{i_{j,k}} a_{B_{\ell;2,k,j}} + R\frac{1+20\delta}{\Gamma} d_{\gamma,\ell} . \tag{3.5}$$

To this end we consider a similar optimization problem for a process that is related to this quantity. Denote by $J = \{j \in [\Gamma] : d_{\gamma,\ell,j} \le \Gamma^{-2/3} d_{\gamma,\ell}\}$. By Cauchy-Schwartz inequality, we see that

$$\sum_{j\in J} d^2_{\gamma,\ell,j} \ge \frac{(1-\Gamma^{-1/5})d^2_{\gamma,\ell}}{\Gamma}\,. \tag{3.6}$$

Write $J = \{j_1, \dots, j_{|J|}\}$ with $1 \le j_1 < j_2 < \cdots < j_{|J|} \le \Gamma$, and define the sequence

$$g_{\gamma,i} = c_\ell^2 \gamma^2 \textstyle\sum_{k\le i} d^2_{\gamma,\ell,j_k} \text{ for } 1 \le i \le 2|J|\,,$$

where $j_i = \Gamma + j_{i-|J|}$ and $d_{\gamma,\ell,j_i} = d_{\gamma,\ell,j_{i-|J|}}$ for $i > |J|$. Consider the process

$$S_{t;\ell} = c_\ell \gamma \textstyle\sum_{0\le k\le i} d_{\gamma,\ell,j_{k+1}} (W^\ell_{t_i\wedge j_{k+1}} - W^\ell_{t_i\wedge(j_{k+1}-1)}) \text{ for } g_{\gamma,i} \le t \le g_{\gamma,i+1}\,,$$

where $t_i = j_{i+1} - 1 + \frac{t-g_{\gamma,i}}{c_\ell^2\gamma^2 d^2_{\gamma,\ell,j_{i+1}}}$ and $g_{2;\gamma,0} = 0$. Thus $\{S_{t;\ell}\}_{0\le t\le g_{\gamma,2|J|}}$ is a standard Brownian motion whose increment between the time points $g_{\gamma,i-1}$ and $g_{\gamma,i}$ is given by $c_\ell\gamma d_{\gamma,\ell,j_i} a_{B_{\ell;2,k,j_i}}$ for $1 \le i \le 2|J|$. Now suppose that $[T_0, T_1, \cdots, T_{R'+1}]$ is the partition of $[0, g_{\gamma,2|J|}]$ given by Theorem 2.3 when applied to $\{S_{t;\ell}\}_{0\le t\le g_{\gamma,2|J|}}$ for the penalty term $\frac{(1+20\delta)d_{\gamma,\ell}}{\Gamma}$ (with appropriate change of scales). Using this we will define L-coarsening of the path that we will construct. Since this has to be simple by our induction hypothesis, it suffices to assign an element in $\{-1,1\}^{2|J|}$ to $j_1, j_2, \cdots, j_{2|J|}$. For this purpose define the random variables $\mathrm{sgn}_{1;\ell}, \cdots, \mathrm{sgn}_{2|J|;\ell}$ as follows:

$$\mathrm{sgn}_{i;\ell} = \begin{cases} -1 & \text{if } T_r < g_{2;\gamma,i} \le T_{r+1} \text{ such that } S_{T_r;\ell} < S_{T_{r+1};\ell} \\ 1 & \text{if } T_r < g_{2;\gamma,i} \le T_{r+1} \text{ such that } S_{T_r;\ell} > S_{T_{r+1};\ell}\,. \end{cases}$$

It is clear that $(\mathrm{sgn}_{1;\ell}, \cdots, \mathrm{sgn}_{2|J|;\ell})$ is identically distributed as $(-\mathrm{sgn}_{1;\ell}, \cdots, -\mathrm{sgn}_{2|J|;\ell})$. As a consequence the distribution of $\pi^{*,\ell+1}_{L,coarse}$ is invariant relative to reflection about the line $y = L - 0.5$.

Now we collect the contributions to the weight of $\pi^{*,\ell+1}$ from all the sources. We use crossings through $V_{\mathrm{up}}, V_{\mathrm{down}}, V_{\mathrm{left}}$ and V_{right}, as in Case 1, to link the crossings $\pi^{*,\ell}_{i,1}$ and $\pi^{*,\ell}_{i,1}$ for $i \in [2]$. As discussed in Case 1, this will not tamper with induction hypothesis 3. We also need a bound on the error we made by interpolating $(S_{t;\ell})$ as a Brownian motion. Notice that for each (T_r, T_{r+1}) that is contained in some interval $(g_{2;\gamma,i-1}, g_{2;\gamma,i})$, we lose at most M_{dis} due to "discretization" of the optimal partition obtained from Theorem 2.3, where

$$M_{\mathrm{dis}} = \max_{1\le i\le 2|J|} \max_{g_{2;\gamma,i-1}\le s,t\le g_{2;\gamma,i}} |S_{t;\ell} - S_{s;\ell}|\,.$$

Thus the total loss we incur due to discretization is at most $M_{\mathrm{dis}}R'$. We use a simple estimate for M_{dis} as follows:

$$\mathbb{E}M_{\mathrm{dis}} \le O(\gamma\Gamma^{-2/3}\log(\Gamma))d_{\gamma,\ell}, \text{ and } \mathbb{P}(|M_{\mathrm{dis}} - \mathbb{E}M_{\mathrm{dis}}| \ge u\gamma\Gamma^{-2/3}\log(\Gamma)d_{\gamma,\ell}) \le e^{-\Omega(u^2)}\,. \tag{3.7}$$

Finally we need to make amends for the fact that what we pass as "penalty" term to (3.5) is actually based on the Gaussians up to level ℓ. But the multiplier at level $\ell+1$ can be at most $\mathrm{e}^{M_{\ell+1}}$, where $M_{\ell+1}$ is the maximum of all Gaussians added at level $\ell+1$. Altogether, we deduce that

$$\begin{aligned} d_{\gamma,\ell+1} \le{}& 2d_{\gamma,\ell}(1+\gamma^2/2 + O(\gamma^3) + \frac{10}{\Gamma}) - \mathcal{I}_\ell + \sqrt{\mathbb{E}R'^2}\sqrt{\mathbb{E}M^2_{\mathrm{dis}}} \\ &+ \frac{(1+20\delta)d_{\gamma,\ell}}{\Gamma}\sqrt{\mathbb{E}R'^2}\sqrt{\mathbb{E}(\mathrm{e}^{\gamma M_{\ell+1}}-1)^2}\,, \end{aligned} \tag{3.8}$$

where

$$\mathcal{I}_\ell = \mathbb{E}\Big(\sum_{i=1}^{R'+1} |S_{T_i;\ell} - S_{T_{i-1};\ell}| - R' \frac{(1+20\delta) d_{\gamma,\ell}}{\Gamma} \Big) ,$$

It follows from condition (d) in Theorem 2.3 and a straightforward computation that

$$\mathbb{E}R'^2 \le O\Big(\Big(\frac{\Gamma^2\gamma^2 \sum_{j\in J} d^2_{\gamma,\ell,j}}{d^2_{\gamma,\ell}} \Big)^2 \Big) . \tag{3.9}$$

In addition, by Theorem 2.3, we deduce that for all $\alpha > 0$

$$\mathcal{I}_\ell \ge (\frac{4}{3} - \delta)\gamma^2\Gamma \frac{\sum_{j\in J} d^2_{\gamma,\ell,j}}{d_{\gamma,\ell}} .$$

Here we used the fact that $\ell \ge 60$. Plugging the bounds from (3.7), (3.9) and Lemma 3.2 into (3.8) we obtain

$$d_{\gamma,\ell+1} \le (2 - 0.6\gamma^2) 2\Gamma (2 - 0.2\gamma^2)^\ell \le 2\Gamma (2 - 0.2\gamma^2)^{\ell+1} .$$

This completes the proof of Theorem 1.1.

References

[1] M. Aizenman, J. T. Chayes, L. Chayes, J. Fröhlich, and L. Russo. On a sharp transition from area law to perimeter law in a system of random surfaces. *Comm. Math. Phys.*, 92(1):19–69, 1983.

[2] A. Auffinger, M. Damron, and J. Hanson. 50 years of first passage percolation. Preprint, available at `http://arxiv.org/abs/1511.03262`.

[3] J. T. Chayes, L. Chayes, and R. Durrett. Connectivity properties of Mandelbrot's percolation process. *Probab. Theory Related Fields*, 77(3):307–324, 1988.

[4] L. Chayes. Aspects of the fractal percolation process. In *Fractal geometry and stochastics (Finsterbergen, 1994)*, volume 37 of *Progr. Probab.*, pages 113–143. Birkhäuser, Basel, 1995.

[5] F. M. Dekking and R. W. J. Meester. On the structure of Mandelbrot's percolation process and other random Cantor sets. *J. Statist. Phys.*, 58(5-6):1109–1126, 1990.

[6] J. Ding and S. Goswami. Liouville first passage percolation: the weight exponent is strictly less than 1 at high temperatures. Preprint, available at `https://arxiv.org/abs/1605.08392`.

[7] J. Ding, R. Roy, and O. Zeitouni. Convergence of the centered maximum of log-correlated gaussian fields. Preprint, available at `http://arxiv.org/abs/1503.04588`.

[8] J. Ding and F. Zhang. Non-universality for first passage percolation on the exponential of log-correlated gaussian fields. Preprint, available at `http://arxiv.org/abs/1506.03293`.

[9] A. Dunlap. Expected regularized total variation of brownian motion. Preprint, available at `http://arxiv.org/abs/1511.03633`.

[10] B. Duplantier and S. Sheffield. Liouville quantum gravity and KPZ. *Invent. Math.*, 185(2):333–393, 2011.

[11] G. R. Grimmett and H. Kesten. Percolation since Saint-Flour. In *Percolation theory at Saint-Flour*, Probab. St.-Flour, pages ix–xxvii. Springer, Heidelberg, 2012.

[12] E. Gwynne, N. Holden, and X. Sun. A distance exponent for liouville quantum gravity. Preprint, available at `http://arxiv.org/abs/1606.01214`.

[13] T. Madaule. Maximum of a log-correlated gaussian field. Preprint, available at `http://arxiv.org/abs/1307.1365`.

[14] J. Miller and S. Sheffield. Liouville quantum gravity and the brownian map III: the conformal structure is determined. Preprint, available at `http://arxiv.org/abs/1608.05391`.

[15] J. Miller and S. Sheffield. Quantum loewner evolution. Preprint, available at `http://arxiv.org/abs/1409.7055`.

[16] P. Mörters and Y. Peres. *Brownian motion*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, Cambridge, 2010. With an appendix by Oded Schramm and Wendelin Werner.

[17] A. M. Polyakov. Quantum geometry of bosonic strings. *Phys. Lett. B*, 103(3):207–210, 1981.

[18] R. Rhodes and V. Vargas. Gaussian multiplicative chaos and applications: a review. *Probab. Surv.*, 11:315–392, 2014.